

Applicants: Graham P. Allaway et al.
Serial No.: 09/904,356
Filed: July 12, 2001
Exhibit 13

BEST AVAILABLE COPY

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 :
C07K 14/705, C12N 15/12

A1

(11) International Publication Number:

WO 96/39437

(43) International Publication Date: 12 December 1996 (12.12.96)

(21) International Application Number:

PCT/US95/07173

(22) International Filing Date:

6 June 1995 (06.06.95)

(71) Applicant (for all designated States except US): HUMAN
GENOME SCIENCES, INC. [US/US]; 9410 Key West
Avenue, Rockville, MD 20850-3338 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): LI, YI [CN/US]; 16125
Howard Landing Drive, Gaithersburg, MD 20878 (US).
RUBEN, Steven, M. [US/US]; 18528 Heritage Hills Drive,
Olney, MD 20832 (US).

(74) Agents: OLSTEIN, Elliot, M.; Carella, Byrne, Bain, Gillilan,
Cecchi, Stewart & Olstein, 6 Becker Farm Road, Roseland,
NJ 07068 (US) et al.

(81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH,
CN, CZ, DE, DK, ES, FI, GB, GE, HU, JP, KE, KG, KP,
KR, KZ, LK, LT, LU, LV, MD, MG, MN, MW, MX, NO,
NZ, PL, PT, RO, RU, SD, SE, SI, SK, TJ, TT, UA, US,
UZ, VN, ARIPO patent (KE, MW, SD, SZ, UG), European
patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU,
MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM,
GA, GN, ML, MR, NE, SN, TD, TG).

Published

With international search report.
With an indication in relation to a deposited
microorganism furnished under Rule 13bis separately
from the description.
Date of receipt by the International Bureau:
1 November 1996 (01.11.96)

(54) Title: HUMAN G-PROTEIN CHEMOKINE RECEPTOR HDGMR10

(57) Abstract

Human G-protein chemokine receptor polypeptides and DNA (RNA) encoding such polypeptides and a procedure for producing such polypeptides by recombinant techniques is disclosed. Also disclosed are methods for utilizing such polypeptides for identifying antagonists and agonists to such polypeptides and methods of using the agonists and antagonists therapeutically to treat conditions related to the underexpression and overexpression of the G-protein chemokine receptor polypeptides, respectively. Also disclosed are diagnostic methods for detecting a mutation in the G-protein chemokine receptor nucleic acid sequences and detecting a level of the soluble form of the receptors in a sample derived from a host.

Applicants: Graham P. Allaway et al.
Serial No.: 09/904,356
Filed: July 12, 2001
Exhibit 13

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

HUMAN G-PROTEIN CHEMOKINE RECEPTOR HDGNR10

This invention relates to newly identified polynucleotides, polypeptides encoded by such polynucleotides, the use of such polynucleotides and polypeptides, as well as the production of such polynucleotides and polypeptides. More particularly, the polypeptide of the present invention is a human 7-transmembrane receptor which has been putatively identified as a chemokine receptor, sometimes hereinafter referred to as "G-Protein Chemokine Receptor" or "HDGNR10". The invention also relates to inhibiting the action of such polypeptides.

It is well established that many medically significant biological processes are mediated by proteins participating in signal transduction pathways that involve G-proteins and/or second messengers, e.g., cAMP (Lefkowitz, Nature, 351:353-354 (1991)). Herein these proteins are referred to as proteins participating in pathways with G-proteins or PPG proteins. Some examples of these proteins include the GPC receptors, such as those for adrenergic agents and dopamine (Kobilka, B.K., et al., PNAS, 84:46-50 (1987); Kobilka, B.K., et al., Science, 238:650-656 (1987); Bunzow, J.R., et al., Nature, 336:783-787 (1988)), G-proteins themselves, effector proteins, e.g., phospholipase C, adenylyl cyclase, and phosphodiesterase, and actuator proteins, e.g., protein

kinase A and protein kinase C (Simon, M.I., et al., Science, 252:802-8 (1991)).

For example, in one form of signal transduction, the effect of hormone binding is activation of an enzyme, adenylate cyclase, inside the cell. Enzyme activation by hormones is dependent on the presence of the nucleotide GTP, and GTP also influences hormone binding. A G-protein connects the hormone receptors to adenylate cyclase. G-protein was shown to exchange GTP for bound GDP when activated by hormone receptors. The GTP-carrying form then binds to an activated adenylate cyclase. Hydrolysis of GTP to GDP, catalyzed by the G-protein itself, returns the G-protein to its basal, inactive form. Thus, the G-protein serves a dual role, as an intermediate that relays the signal from receptor to effector, and as a clock that controls the duration of the signal.

The membrane protein gene superfamily of G-protein coupled receptors has been characterized as having seven putative transmembrane domains. The domains are believed to represent transmembrane α -helices connected by extracellular or cytoplasmic loops. G-protein coupled receptors include a wide range of biologically active receptors, such as hormone, viral, growth factor and neuroreceptors.

G-protein coupled receptors have been characterized as including these seven conserved hydrophobic stretches of about 20 to 30 amino acids, connecting at least eight divergent hydrophilic loops. The G-protein family of coupled receptors includes dopamine receptors which bind to neuroleptic drugs used for treating psychotic and neurological disorders. Other examples of members of this family include calcitonin, adrenergic, endothelin, cAMP, adenosine, muscarinic, acetylcholine, serotonin, histamine, thrombin, kinin, follicle stimulating hormone, opsins, endothelial differentiation gene-1 receptor and rhodopsins, odorant, cytomegalovirus receptors, etc.

G-protein coupled receptors can be intracellularly coupled by heterotrimeric G-proteins to various intracellular enzymes, ion channels and transporters (see, Johnson et al., Endoc., Rev., 10:317-331 (1989)). Different G-protein α -subunits preferentially stimulate particular effectors to modulate various biological functions in a cell. Phosphorylation of cytoplasmic residues of G-protein coupled receptors have been identified as an important mechanism for the regulation of G-protein coupling of some G-protein coupled receptors. G-protein coupled receptors are found in numerous sites within a mammalian host.

Chemokines, also referred to as intercrine cytokines, are a subfamily of structurally and functionally related cytokines. These molecules are 8-10 kd in size. In general, chemokines exhibit 20% to 75% homology at the amino acid level and are characterized by four conserved cysteine residues that form two disulfide bonds. Based on the arrangement of the first two cysteine residues, chemokines have been classified into two subfamilies, alpha and beta. In the alpha subfamily, the first two cysteines are separated by one amino acid and hence are referred to as the "C-X-C" subfamily. In the beta subfamily, the two cysteines are in an adjacent position and are, therefore, referred to as the "C-C" subfamily. Thus far, at least nine different members of this family have been identified in humans.

The intercrine cytokines exhibit a wide variety of functions. A hallmark feature is their ability to elicit chemotactic migration of distinct cell types, including monocytes, neutrophils, T lymphocytes, basophils and fibroblasts. Many chemokines have proinflammatory activity and are involved in multiple steps during an inflammatory reaction. These activities include stimulation of histamine release, lysosomal enzyme and leukotriene release, increased adherence of target immune cells to endothelial cells, enhanced binding of complement proteins, induced expression

of granulocyte adhesion molecules and complement receptors, and respiratory burst. In addition to their involvement in inflammation, certain chemokines have been shown to exhibit other activities. For example, macrophage inflammatory protein 1 (MIP-1) is able to suppress hematopoietic stem cell proliferation, platelet factor-4 (PF-4) is a potent inhibitor of endothelial cell growth, Interleukin-8 (IL-8) promotes proliferation of keratinocytes, and GRO is an autocrine growth factor for melanoma cells.

In light of the diverse biological activities, it is not surprising that chemokines have been implicated in a number of physiological and disease conditions, including lymphocyte trafficking, wound healing, hematopoietic regulation and immunological disorders such as allergy, asthma and arthritis.

In accordance with one aspect of the present invention, there are provided novel mature receptor polypeptides as well as biologically active and diagnostically or therapeutically useful fragments, analogs and derivatives thereof. The receptor polypeptides of the present invention are of human origin.

In accordance with another aspect of the present invention, there are provided isolated nucleic acid molecules encoding the receptor polypeptides of the present invention, including mRNAs, DNAs, cDNAs, genomic DNA as well as antisense analogs thereof and biologically active and diagnostically or therapeutically useful fragments thereof.

In accordance with a further aspect of the present invention, there are provided processes for producing such receptor polypeptides by recombinant techniques comprising culturing recombinant prokaryotic and/or eukaryotic host cells, containing nucleic acid sequences encoding the receptor polypeptides of the present invention, under conditions promoting expression of said polypeptides and subsequent recovery of said polypeptides.

In accordance with yet a further aspect of the present invention, there are provided antibodies against such receptor polypeptides.

In accordance with another aspect of the present invention there are provided methods of screening for compounds which bind to and activate or inhibit activation of the receptor polypeptides of the present invention.

In accordance with still another embodiment of the present invention there are provided processes of administering compounds to a host which bind to and activate the receptor polypeptide of the present invention which are useful in stimulating haematopoiesis, wound healing, coagulation, angiogenesis, to treat solid tumors, chronic infections, leukemia, T-cell mediated auto-immune diseases, parasitic infections, psoriasis, and to stimulate growth factor activity.

In accordance with another aspect of the present invention there is provided a method of administering the receptor polypeptides of the present invention via gene therapy to treat conditions related to underexpression of the polypeptides or underexpression of a ligand for the receptor polypeptide.

In accordance with still another embodiment of the present invention there are provided processes of administering compounds to a host which bind to and inhibit activation of the receptor polypeptides of the present invention which are useful in the prevention and/or treatment of allergy, atherogenesis, anaphylaxis, malignancy, chronic and acute inflammation, histamine and IgE-mediated allergic reactions, prostaglandin-independent fever, bone marrow failure, silicosis, sarcoidosis, rheumatoid arthritis, shock and hyper-eosinophilic syndrome.

In accordance with yet another aspect of the present invention, there are provided nucleic acid probes comprising nucleic acid molecules of sufficient length to specifically

hybridize to the polynucleotide sequences of the present invention.

In accordance with still another aspect of the present invention, there are provided diagnostic assays for detecting diseases related to mutations in the nucleic acid sequences encoding such polypeptides and for detecting an altered level of the soluble form of the receptor polypeptides.

In accordance with yet a further aspect of the present invention, there are provided processes for utilizing such receptor polypeptides, or polynucleotides encoding such polypeptides, for in vitro purposes related to scientific research, synthesis of DNA and manufacture of DNA vectors.

These and other aspects of the present invention should be apparent to those skilled in the art from the teachings herein.

The following drawings are illustrative of embodiments of the invention and are not meant to limit the scope of the invention as encompassed by the claims.

Figure 1 shows the cDNA sequence and the corresponding deduced amino acid sequence of the G-protein coupled receptor of the present invention. The standard one-letter abbreviation for amino acids is used. Sequencing was performed using a 373 Automated DNA sequencer (Applied Biosystems, Inc.).

Figure 2 illustrates an amino acid alignment of the G-protein chemokine receptor of the present invention and the human MCP-1 receptor.

In accordance with an aspect of the present invention, there is provided an isolated nucleic acid (polynucleotide) which encodes for the mature polypeptide having the deduced amino acid sequence of Figure 1 (SEQ ID NO:2) or for the mature polypeptide encoded by the cDNA of the clone deposited as ATCC Deposit No. _____ on June 1, 1995.

The polynucleotide of this invention was discovered in a cDNA library derived from human monocytes. It is

structurally related to the G protein-coupled receptor family. It contains an open reading frame encoding a protein of 352 amino acid residues. The protein exhibits the highest degree of homology to a human MCP-1 receptor with 70.1 % identity and 82.9 % similarity over a 347 amino acid stretch.

The polynucleotide of the present invention may be in the form of RNA or in the form of DNA, which DNA includes cDNA, genomic DNA, and synthetic DNA. The DNA may be double-stranded or single-stranded, and if single stranded may be the coding strand or non-coding (anti-sense) strand. The coding sequence which encodes the mature polypeptide may be identical to the coding sequence shown in Figure 1 (SEQ ID NO:1) or that of the deposited clone or may be a different coding sequence which coding sequence, as a result of the redundancy or degeneracy of the genetic code, encodes the same mature polypeptide as the DNA of Figure 1 (SEQ ID NO:1) or the deposited cDNA.

The polynucleotide which encodes for the mature polypeptide of Figure 1 or for the mature polypeptide encoded by the deposited cDNA may include: only the coding sequence for the mature polypeptide; the coding sequence for the mature polypeptide and additional coding sequence such as a transmembrane (TM) or intra-cellular domain; the coding sequence for the mature polypeptide (and optionally additional coding sequence) and non-coding sequence, such as introns or non-coding sequence 5' and/or 3' of the coding sequence for the mature polypeptide.

Thus, the term "polynucleotide encoding a polypeptide" encompasses a polynucleotide which includes only coding sequence for the polypeptide as well as a polynucleotide which includes additional coding and/or non-coding sequence.

The present invention further relates to variants of the hereinabove described polynucleotides which encode for fragments, analogs and derivatives of the polypeptide having the deduced amino acid sequence of Figure 1 or the

polypeptide encoded by the cDNA of the deposited clone. The variant of the polynucleotide may be a naturally occurring allelic variant of the polynucleotide or a non-naturally occurring variant of the polynucleotide. ✓

Thus, the present invention includes polynucleotides encoding the same mature polypeptide as shown in Figure 1 (SEQ ID NO:2) or the same mature polypeptide encoded by the cDNA of the deposited clone as well as variants of such polynucleotides which variants encode for a fragment, derivative or analog of the polypeptide of Figure 1 (SEQ ID NO:2) or the polypeptide encoded by the cDNA of the deposited clone. Such nucleotide variants include deletion variants, substitution variants and addition or insertion variants. ✓

As hereinabove indicated, the polynucleotide may have a coding sequence which is a naturally occurring allelic variant of the coding sequence shown in Figure 1 (SEQ ID NO:1) or of the coding sequence of the deposited clone. As known in the art, an allelic variant is an alternate form of a polynucleotide sequence which may have a substitution, deletion or addition of one or more nucleotides, which does not substantially alter the function of the encoded polypeptide. ✓

The polynucleotides may also encode for a soluble form of the G-protein chemokine receptor polypeptide which is the extracellular portion of the polypeptide which has been cleaved from the TM and intracellular domain of the full-length polypeptide of the present invention. ✓

The polynucleotides of the present invention may also have the coding sequence fused in frame to a marker sequence which allows for purification of the polypeptide of the present invention. The marker sequence may be a hexahistidine tag supplied by a pQE-9 vector to provide for purification of the mature polypeptide fused to the marker in the case of a bacterial host, or, for example, the marker sequence may be a hemagglutinin (HA) tag when a mammalian

✓ host, e.g. COS-7 cells, is used. The HA tag corresponds to an epitope derived from the influenza hemagglutinin protein (Wilson, I., et al., Cell, 37:767 (1984)).

The term "gene" means the segment of DNA involved in producing a polypeptide chain; it includes regions preceding and following the coding region (leader and trailer) as well as intervening sequences (introns) between individual coding segments (exons).

Fragments of the full length gene of the present invention may be used as a hybridization probe for a cDNA library to isolate the full length cDNA and to isolate other cDNAs which have a high sequence similarity to the gene or similar biological activity. Probes of this type preferably have at least 30 bases and may contain, for example, 50 or more bases. The probe may also be used to identify a cDNA clone corresponding to a full length transcript and a genomic clone or clones that contain the complete gene including regulatory and promotor regions, exons, and introns. An example of a screen comprises isolating the coding region of the gene by using the known DNA sequence to synthesize an oligonucleotide probe. Labeled oligonucleotides having a sequence complementary to that of the gene of the present invention are used to screen a library of human cDNA, genomic DNA or mRNA to determine which members of the library the probe hybridizes to.

The present invention further relates to polynucleotides which hybridize to the hereinabove-described sequences if there is at least 70%, preferably at least 90%, and more preferably at least 95% identity between the sequences. The present invention particularly relates to polynucleotides which hybridize under stringent conditions to the hereinabove-described polynucleotides. As herein used, the term "stringent conditions" means hybridization will occur only if there is at least 95% and preferably at least 97% identity between the sequences. The polynucleotides

which hybridize to the hereinabove described polynucleotides in a preferred embodiment encode polypeptides which either retain substantially the same biological function or activity as the mature polypeptide encoded by the cDNAs of Figure 1 (SEQ ID NO:1) or the deposited cDNA(s).

Alternatively, the polynucleotide may have at least 20 bases, preferably 30 bases, and more preferably at least 50 bases which hybridize to a polynucleotide of the present invention and which has an identity thereto, as hereinabove described, and which may or may not retain activity. For example, such polynucleotides may be employed as probes for the polynucleotide of SEQ ID NO:1, for example, for recovery of the polynucleotide or as a diagnostic probe or as a PCR primer.

Thus, the present invention is directed to polynucleotides having at least a 70% identity, preferably at least 90% and more preferably at least a 95% identity to a polynucleotide which encodes the polypeptide of SEQ ID NO:2 as well as fragments thereof, which fragments have at least 30 bases and preferably at least 50 bases and to polypeptides encoded by such polynucleotides.

The deposit(s) referred to herein will be maintained under the terms of the Budapest Treaty on the International Recognition of the Deposit of Micro-organisms for purposes of Patent Procedure. These deposits are provided merely as convenience to those of skill in the art and are not an admission that a deposit is required under 35 U.S.C. §112. The sequence of the polynucleotides contained in the deposited materials, as well as the amino acid sequence of the polypeptides encoded thereby, are incorporated herein by reference and are controlling in the event of any conflict with any description of sequences herein. A license may be required to make, or sell the deposited materials, and no such license is hereby granted.

The present invention further relates to a G-protein chemokine receptor polypeptide which has the deduced amino acid sequence of Figure 1 (SEQ ID NO:2) or which has the amino acid sequence encoded by the deposited cDNA, as well as fragments, analogs and derivatives of such polypeptide.

The terms "fragment," "derivative" and "analog" when referring to the polypeptide of Figure 1 or that encoded by the deposited cDNA, means a polypeptide which either retains substantially the same biological function or activity as such polypeptide, i.e. functions as a G-protein chemokine receptor, or retains the ability to bind the ligand or the receptor even though the polypeptide does not function as a G-protein chemokine receptor, for example, a soluble form of the receptor. An analog includes a proprotein which can be activated by cleavage of the proprotein portion to produce an active mature polypeptide.

The polypeptide of the present invention may be a recombinant polypeptide, a natural polypeptide or a synthetic polypeptide, preferably a recombinant polypeptide.

The fragment, derivative or analog of the polypeptide of Figure 1 (SEQ ID NO:2) or that encoded by the deposited cDNA may be (i) one in which one or more of the amino acid residues are substituted with a conserved or non-conserved amino acid residue (preferably a conserved amino acid residue) and such substituted amino acid residue may or may not be one encoded by the genetic code, or (ii) one in which one or more of the amino acid residues includes a substituent group, or (iii) one in which the mature polypeptide is fused with another compound, such as a compound to increase the half-life of the polypeptide (for example, polyethylene glycol), or (iv) one in which the additional amino acids are fused to the mature polypeptide for purification of the polypeptide or (v) one in which a fragment of the polypeptide is soluble, i.e. not membrane bound, yet still binds ligands to the membrane bound receptor. Such fragments, derivatives

and analogs are deemed to be within the scope of those skilled in the art from the teachings herein. ✓

The polypeptides and polynucleotides of the present invention are preferably provided in an isolated form, and preferably are purified to homogeneity. ✓

The polypeptides of the present invention include the polypeptide of SEQ ID NO:2 (in particular the mature polypeptide) as well as polypeptides which have at least 70% similarity (preferably a 70% identity) to the polypeptide of SEQ ID NO:2 and more preferably a 90% similarity (more preferably a 90% identity) to the polypeptide of SEQ ID NO:2 and still more preferably a 95% similarity (still more preferably a 90% identity) to the polypeptide of SEQ ID NO:2 and to portions of such polypeptide with such portion of the polypeptide generally containing at least 30 amino acids and more preferably at least 50 amino acids. 2.

As known in the art "similarity" between two polypeptides is determined by comparing the amino acid sequence and conserved amino acid substitutes thereto of the polypeptide to the sequence of a second polypeptide. ✓

Fragments or portions of the polypeptides of the present invention may be employed for producing the corresponding full-length polypeptide by peptide synthesis, therefore, the fragments may be employed as intermediates for producing the full-length polypeptides. ✓
Fragments or portions of the polynucleotides of the present invention may be used to synthesize full-length polynucleotides of the present invention.

The term "gene" means the segment of DNA involved in producing a polypeptide chain; it includes regions preceding and following the coding region "leader and trailer" as well as intervening sequences (introns) between individual coding segments (exons). ✓

The term "isolated" means that the material is removed from its original environment (e.g., the natural environment

if it is naturally occurring). For example, a naturally-occurring polynucleotide or polypeptide present in a living animal is not isolated, but the same polynucleotide or polypeptide, separated from some or all of the coexisting materials in the natural system, is isolated. Such polynucleotides could be part of a vector and/or such polynucleotides or polypeptides could be part of a composition, and still be isolated in that such vector or composition is not part of its natural environment. ✓

The polypeptides of the present invention include the polypeptide of SEQ ID NO:2 (in particular the mature polypeptide) as well as polypeptides which have at least 70% similarity (preferably at least 70% identity) to the polypeptide of SEQ ID NO:2 and more preferably at least 90% similarity (more preferably at least 90% identity) to the polypeptide of SEQ ID NO:2 and still more preferably at least 95% similarity (still more preferably at least 95% identity) to the polypeptide of SEQ ID NO:2 and also include portions of such polypeptides with such portion of the polypeptide generally containing at least 30 amino acids and more preferably at least 50 amino acids. ✓

As known in the art "similarity" between two polypeptides is determined by comparing the amino acid sequence and its conserved amino acid substitutes of one polypeptide to the sequence of a second polypeptide. ✓

Fragments or portions of the polypeptides of the present invention may be employed for producing the corresponding full-length polypeptide by peptide synthesis; therefore, the fragments may be employed as intermediates for producing the full-length polypeptides. ✓
Fragments or portions of the polynucleotides of the present invention may be used to synthesize full-length polynucleotides of the present invention.

The present invention also relates to vectors which include polynucleotides of the present invention, host cells

which are genetically engineered with vectors of the invention and the production of polypeptides of the invention by recombinant techniques.

Host cells are genetically engineered (transduced or transformed or transfected) with the vectors of this invention which may be, for example, a cloning vector or an expression vector. The vector may be, for example, in the form of a plasmid, a viral particle, a phage, etc. The engineered host cells can be cultured in conventional nutrient media modified as appropriate for activating promoters, selecting transformants or amplifying the genes of the present invention. The culture conditions, such as temperature, pH and the like, are those previously used with the host cell selected for expression, and will be apparent to the ordinarily skilled artisan.

The polynucleotides of the present invention may be employed for producing polypeptides by recombinant techniques. Thus, for example, the polynucleotide may be included in any one of a variety of expression vectors for expressing a polypeptide. Such vectors include chromosomal, nonchromosomal and synthetic DNA sequences, e.g., derivatives of SV40; bacterial plasmids; phage DNA; baculovirus; yeast plasmids; vectors derived from combinations of plasmids and phage DNA, viral DNA such as vaccinia, adenovirus, fowl pox virus, and pseudorabies. However, any other vector may be used as long as it is replicable and viable in the host.

The appropriate DNA sequence may be inserted into the vector by a variety of procedures. In general, the DNA sequence is inserted into an appropriate restriction endonuclease site(s) by procedures known in the art. Such procedures and others are deemed to be within the scope of those skilled in the art.

The DNA sequence in the expression vector is operatively linked to an appropriate expression control sequence(s)

(promoter) to direct mRNA synthesis. As representative examples of such promoters, there may be mentioned: LTR or SV40 promoter, the E. coli lac or trp, the phage lambda P_L promoter and other promoters known to control expression of genes in prokaryotic or eukaryotic cells or their viruses. The expression vector also contains a ribosome binding site for translation initiation and a transcription terminator. The vector may also include appropriate sequences for amplifying expression.

In addition, the expression vectors preferably contain one or more selectable marker genes to provide a phenotypic trait for selection of transformed host cells such as dihydrofolate reductase or neomycin resistance for eukaryotic cell culture, or such as tetracycline or ampicillin resistance in E. coli.

The vector containing the appropriate DNA sequence as hereinabove described, as well as an appropriate promoter or control sequence, may be employed to transform an appropriate host to permit the host to express the protein.

As representative examples of appropriate hosts, there may be mentioned: bacterial cells, such as E. coli, Streptomyces, Salmonella typhimurium; fungal cells, such as yeast; insect cells such as Drosophila and Spodoptera Sf9; animal cells such as CHO, COS or Bowes melanoma; adenovirus; plant cells, etc. The selection of an appropriate host is deemed to be within the scope of those skilled in the art from the teachings herein.

More particularly, the present invention also includes recombinant constructs comprising one or more of the sequences as broadly described above. The constructs comprise a vector, such as a plasmid or viral vector, into which a sequence of the invention has been inserted, in a forward or reverse orientation. In a preferred aspect of this embodiment, the construct further comprises regulatory sequences, including, for example, a promoter, operably

linked to the sequence. Large numbers of suitable vectors and promoters are known to those of skill in the art, and are commercially available. The following vectors are provided by way of example. Bacterial: pQE70, pQE60, pQE-9 (Qiagen), pbs, pD10, phagescript, psiX174, pbluescript SK, pbsks, pNH8A, pNH16a, pNH18A, pNH46A (Stratagene); ptrc99a, pKK223-3, pKK233-3, pDR540, pRIT5 (Pharmacia). Eukaryotic: pWLNBO, pSV2CAT, pOG44, pXT1, pSG (Stratagene) pSVK3, pBPV, pMSG, pSVL (Pharmacia). However, any other plasmid or vector may be used as long as they are replicable and viable in the host.

Promoter regions can be selected from any desired gene using CAT (chloramphenicol transferase) vectors or other vectors with selectable markers. Two appropriate vectors are pKK232-8 and PCM7. Particular named bacterial promoters include lacI, lacZ, T3, T7, gpt, lambda P_R, P_L and trp. Eukaryotic promoters include CMV immediate early, HSV thymidine kinase, early and late SV40, LTRs from retrovirus, and mouse metallothionein-I. Selection of the appropriate vector and promoter is well within the level of ordinary skill in the art.

In a further embodiment, the present invention relates to host cells containing the above-described constructs. The host cell can be a higher eukaryotic cell, such as a mammalian cell, or a lower eukaryotic cell, such as a yeast cell, or the host cell can be a prokaryotic cell, such as a bacterial cell. Introduction of the construct into the host cell can be effected by calcium phosphate transfection, DEAE-Dextran mediated transfection, or electroporation. (Davis, L., Dibner, M., Battey, I., Basic Methods in Molecular Biology, (1986)).

The constructs in host cells can be used in a conventional manner to produce the gene product encoded by the recombinant sequence. Alternatively, the polypeptides of

the invention can be synthetically produced by conventional peptide synthesizers.

Mature proteins can be expressed in mammalian cells, yeast, bacteria, or other cells under the control of appropriate promoters. Cell-free translation systems can also be employed to produce such proteins using RNAs derived from the DNA constructs of the present invention. Appropriate cloning and expression vectors for use with prokaryotic and eukaryotic hosts are described by Sambrook, et al., *Molecular Cloning: A Laboratory Manual*, Second Edition, Cold Spring Harbor, N.Y., (1989), the disclosure of which is hereby incorporated by reference.

Transcription of the DNA encoding the polypeptides of the present invention by higher eukaryotes is increased by inserting an enhancer sequence into the vector. Enhancers are cis-acting elements of DNA, usually about from 10 to 300 bp that act on a promoter to increase its transcription. Examples including the SV40 enhancer on the late side of the replication origin bp 100 to 270, a cytomegalovirus early promoter enhancer, the polyoma enhancer on the late side of the replication origin, and adenovirus enhancers.

Generally, recombinant expression vectors will include origins of replication and selectable markers permitting transformation of the host cell, e.g., the ampicillin resistance gene of E. coli and S. cerevisiae TRP1 gene, and a promoter derived from a highly-expressed gene to direct transcription of a downstream structural sequence. Such promoters can be derived from operons encoding glycolytic enzymes such as 3-phosphoglycerate kinase (PGK), α -factor, acid phosphatase, or heat shock proteins, among others. The heterologous structural sequence is assembled in appropriate phase with translation initiation and termination sequences, and preferably, a leader sequence capable of directing secretion of translated protein into the periplasmic space or extracellular medium. Optionally, the heterologous sequence

can encode a fusion protein including an N-terminal identification peptide imparting desired characteristics, e.g., stabilization or simplified purification of expressed recombinant product.

Useful expression vectors for bacterial use are constructed by inserting a structural DNA sequence encoding a desired protein together with suitable translation initiation and termination signals in operable reading phase with a functional promoter. The vector will comprise one or more phenotypic selectable markers and an origin of replication to ensure maintenance of the vector and to, if desirable, provide amplification within the host. Suitable prokaryotic hosts for transformation include E. coli, Bacillus subtilis, Salmonella typhimurium and various species within the genera Pseudomonas, Streptomyces, and Staphylococcus, although others may also be employed as a matter of choice.

As a representative but nonlimiting example, useful expression vectors for bacterial use can comprise a selectable marker and bacterial origin of replication derived from commercially available plasmids comprising genetic elements of the well known cloning vector pBR322 (ATCC 37017). Such commercial vectors include, for example, pKK223-3 (Pharmacia Fine Chemicals, Uppsala, Sweden) and GEM1 (Promega Biotec, Madison, WI, USA). These pBR322 "backbone" sections are combined with an appropriate promoter and the structural sequence to be expressed.

Following transformation of a suitable host strain and growth of the host strain to an appropriate cell density, the selected promoter is induced by appropriate means (e.g., temperature shift or chemical induction) and cells are cultured for an additional period.

Cells are typically harvested by centrifugation, disrupted by physical or chemical means, and the resulting crude extract retained for further purification.

Microbial cells employed in expression of proteins can be disrupted by any convenient method, including freeze-thaw cycling, sonication, mechanical disruption, or use of cell lysing agents, such methods are well know to those skilled in the art.

Various mammalian cell culture systems can also be employed to express recombinant protein. Examples of mammalian expression systems include the COS-7 lines of monkey kidney fibroblasts, described by Gluzman, Cell, 23:175 (1981), and other cell lines capable of expressing a compatible vector, for example, the C127, 3T3, CHO, HeLa and BHK cell lines. Mammalian expression vectors will comprise an origin of replication, a suitable promoter and enhancer, and also any necessary ribosome binding sites, polyadenylation site, splice donor and acceptor sites, transcriptional termination sequences, and 5' flanking nontranscribed sequences. DNA sequences derived from the SV40 splice, and polyadenylation sites may be used to provide the required nontranscribed genetic elements.

The G-protein chemokine receptor polypeptides can be recovered and purified from recombinant cell cultures by methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Protein refolding steps can be used, as necessary, in completing configuration of the mature protein. Finally, high performance liquid chromatography (HPLC) can be employed for final purification steps.

The polypeptides of the present invention may be a naturally purified product, or a product of chemical synthetic procedures, or produced by recombinant techniques from a prokaryotic or eukaryotic host (for example, by bacterial, yeast, higher plant, insect and mammalian cells in

culture). Depending upon the host employed in a recombinant production procedure, the polypeptides of the present invention may be glycosylated or may be non-glycosylated. Polypeptides of the invention may also include an initial methionine amino acid residue. ✓

The polynucleotides and polypeptides of the present invention may be employed as research reagents and materials for discovery of treatments and diagnostics to human disease. ✓

The G-protein chemokine receptors of the present invention may be employed in a process for screening for compounds which activate (agonists) or inhibit activation (antagonists) of the receptor polypeptide of the present invention. ✓

In general, such screening procedures involve providing appropriate cells which express the receptor polypeptide of the present invention on the surface thereof. Such cells include cells from mammals, yeast, drosophila or *E. Coli*. In particular, a polynucleotide encoding the receptor of the present invention is employed to transfect cells to thereby express the G-protein chemokine receptor. The expressed receptor is then contacted with a test compound to observe binding, stimulation or inhibition of a functional response. ✓

One such screening procedure involves the use of melanophores which are transfected to express the G-protein chemokine receptor of the present invention. Such a screening technique is described in PCT WO 92/01810 published February 6, 1992. ✓

Thus, for example, such assay may be employed for screening for a compound which inhibits activation of the receptor polypeptide of the present invention by contacting the melanophore cells which encode the receptor with both the receptor ligand and a compound to be screened. Inhibition of the signal generated by the ligand indicates that a compound is a potential antagonist for the receptor, i.e., inhibits activation of the receptor. ✓

The screen may be employed for determining a compound which activates the receptor by contacting such cells with compounds to be screened and determining whether such compound generates a signal, i.e., activates the receptor. ✓

Other screening techniques include the use of cells which express the G-protein chemokine receptor (for example, transfected CHO cells) in a system which measures extracellular pH changes caused by receptor activation, for example, as described in Science, volume 246, pages 181-296 (October 1989). For example, compounds may be contacted with a cell which expresses the receptor polypeptide of the present invention and a second messenger response, e.g. signal transduction or pH changes, may be measured to determine whether the potential compound activates or inhibits the receptor. ✓

Another such screening technique involves introducing RNA encoding the G-protein chemokine receptor into *Xenopus* oocytes to transiently express the receptor. The receptor oocytes may then be contacted with the receptor ligand and a compound to be screened, followed by detection of inhibition or activation of a calcium signal in the case of screening for compounds which are thought to inhibit activation of the receptor. ✓

Another screening technique involves expressing the G-protein chemokine receptor in which the receptor is linked to a phospholipase C or D. As representative examples of such cells, there may be mentioned endothelial cells, smooth muscle cells, embryonic kidney cells, etc. The screening may be accomplished as hereinabove described by detecting activation of the receptor or inhibition of activation of the receptor from the phospholipase second signal. ✓

Another method involves screening for compounds which inhibit activation of the receptor polypeptide of the present invention antagonists by determining inhibition binding of labeled ligand to cells which have the receptor on the

surface thereof. Such a method involves transfecting a eukaryotic cell with DNA encoding the G-protein chemokine receptor such that the cell expresses the receptor on its surface and contacting the cell with a compound in the presence of a labeled form of a known ligand. The ligand can be labeled, e.g., by radioactivity. The amount of labeled ligand bound to the receptors is measured, e.g., by measuring radioactivity of the receptors. If the compound binds to the receptor as determined by a reduction of labeled ligand which binds to the receptors, the binding of labeled ligand to the receptor is inhibited.

An antibody may antagonize a G-protein chemokine receptor of the present invention, or in some cases an oligopeptide, which bind to the G-protein chemokine receptor but does not elicit a second messenger response such that the activity of the G-protein chemokine receptors is prevented. Antibodies include anti-idiotypic antibodies which recognize unique determinants generally associated with the antigen-binding site of an antibody. Potential antagonist compounds also include proteins which are closely related to the ligand of the G-protein chemokine receptors, i.e. a fragment of the ligand, which have lost biological function and when binding to the G-protein chemokine receptor elicit no response.

An antisense construct prepared through the use of antisense technology, may be used to control gene expression through triple-helix formation or antisense DNA or RNA, both of which methods are based on binding of a polynucleotide to DNA or RNA. For example, the 5' coding portion of the polynucleotide sequence, which encodes for the mature polypeptides of the present invention, is used to design an antisense RNA oligonucleotide of from about 10 to 40 base pairs in length. A DNA oligonucleotide is designed to be complementary to a region of the gene involved in transcription (triple helix -see Lee et al., Nucl. Acids Res., 6:3073 (1979); Cooney et al, Science, 241:456 (1988);

and Dervan et al., Science, 251: 1360 (1991)), thereby preventing transcription and the production of G-protein chemokine receptor. The antisense RNA oligonucleotide hybridizes to the mRNA *in vivo* and blocks translation of mRNA molecules into G-protein coupled receptor (antisense - Okano, J. Neurochem., 56:560 (1991); Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression, CRC Press, Boca Raton, FL (1988)). The oligonucleotides described above can also be delivered to cells such that the antisense RNA or DNA may be expressed *in vivo* to inhibit production of G-protein chemokine receptor. ✓

A small molecule which binds to the G-protein chemokine receptor, making it inaccessible to ligands such that normal biological activity is prevented, for example small peptides or peptide-like molecules, may also be used to inhibit activation of the receptor polypeptide of the present invention. ✓

A soluble form of the G-protein chemokine receptor, e.g. a fragment of the receptors, may be used to inhibit activation of the receptor by binding to the ligand to a polypeptide of the present invention and preventing the ligand from interacting with membrane bound G-protein chemokine receptors. ✓

The compounds which bind to and activate the G-protein chemokine receptors of the present invention may be employed to stimulate haematopoiesis, wound healing, coagulation, angiogenesis, to treat solid tumors, chronic infections, leukemia, T-cell mediated auto-immune diseases, parasitic infections, psoriasis, and to stimulate growth factor activity. ✓

The compounds which bind to and inhibit the G-protein chemokine receptors of the present invention may be employed to treat allergy, atherogenesis, anaphylaxis, malignancy, chronic and acute inflammation, histamine and IgE-mediated allergic reactions, prostaglandin-independent fever, bone

marrow failure, silicosis, sarcoidosis, rheumatoid arthritis, shock and hyper-eosinophilic syndrome.

The compounds may be employed in combination with a suitable pharmaceutical carrier. Such compositions comprise a therapeutically effective amount of the compound and a pharmaceutically acceptable carrier or excipient. Such a carrier includes but is not limited to saline, buffered saline, dextrose, water, glycerol, ethanol, and combinations thereof. The formulation should suit the mode of administration.

The invention also provides a pharmaceutical pack or kit comprising one or more containers filled with one or more of the ingredients of the pharmaceutical compositions of the invention. Associated with such container(s) can be a notice in the form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or biological products, which notice reflects approval by the agency of manufacture, use or sale for human administration. In addition, the compounds of the present invention may be employed in conjunction with other therapeutic compounds.

The pharmaceutical compositions may be administered in a convenient manner such as by the topical, intravenous, intraperitoneal, intramuscular, subcutaneous, intranasal or intradermal (applicable?) routes. The pharmaceutical compositions are administered in an amount which is effective for treating and/or prophylaxis of the specific indication. In general, the pharmaceutical compositions will be administered in an amount of at least about 10 μ g/kg body weight and in most cases they will be administered in an amount not in excess of about 8 mg/Kg body weight per day. In most cases, the dosage is from about 10 μ g/kg to about 1 mg/kg body weight daily, taking into account the routes of administration, symptoms, etc. (CONFIRM DOSAGES)

The G-protein chemokine receptor polypeptides and antagonists or agonists which are polypeptides, may also be

employed in accordance with the present invention by expression of such polypeptides *in vivo*, which is often referred to as "gene therapy."

Thus, for example, cells from a patient may be engineered with a polynucleotide (DNA or RNA) encoding a polypeptide *ex vivo*, with the engineered cells then being provided to a patient to be treated with the polypeptide. Such methods are well-known in the art. For example, cells may be engineered by procedures known in the art by use of a retroviral particle containing RNA encoding a polypeptide of the present invention.

Similarly, cells may be engineered *in vivo* for expression of a polypeptide *in vivo* by, for example, procedures known in the art. As known in the art, a producer cell for producing a retroviral particle containing RNA encoding the polypeptide of the present invention may be administered to a patient for engineering cells *in vivo* and expression of the polypeptide *in vivo*. These and other methods for administering a polypeptide of the present invention by such method should be apparent to those skilled in the art from the teachings of the present invention. For example, the expression vehicle for engineering cells may be other than a retrovirus, for example, an adenovirus which may be used to engineer cells *in vivo* after combination with a suitable delivery vehicle.

Retroviruses from which the retroviral plasmid vectors hereinabove mentioned may be derived include, but are not limited to, Moloney Murine Leukemia Virus, spleen necrosis virus, retroviruses such as Rous Sarcoma Virus, Harvey Sarcoma Virus, avian leukosis virus, gibbon ape leukemia virus, human immunodeficiency virus, adenovirus, Myeloproliferative Sarcoma Virus, and mammary tumor virus. In one embodiment, the retroviral plasmid vector is derived from Moloney Murine Leukemia Virus.

The vector includes one or more promoters. Suitable promoters which may be employed include, but are not limited to, the retroviral LTR; the SV40 promoter; and the human cytomegalovirus (CMV) promoter described in Miller, et al., Biotechniques, Vol. 7, No. 9, 980-990 (1989), or any other promoter (e.g., cellular promoters such as eukaryotic cellular promoters including, but not limited to, the histone, pol III, and β -actin promoters). Other viral promoters which may be employed include, but are not limited to, adenovirus promoters, thymidine kinase (TK) promoters, and B19 parvovirus promoters. The selection of a suitable promoter will be apparent to those skilled in the art from the teachings contained herein.

The nucleic acid sequence encoding the polypeptide of the present invention is under the control of a suitable promoter. Suitable promoters which may be employed include, but are not limited to, adenoviral promoters, such as the adenoviral major late promoter; or heterologous promoters, such as the cytomegalovirus (CMV) promoter; the respiratory syncytial virus (RSV) promoter; inducible promoters, such as the MMT promoter, the metallothionein promoter; heat shock promoters; the albumin promoter; the ApoA1 promoter; human globin promoters; viral thymidine kinase promoters, such as the Herpes Simplex thymidine kinase promoter; retroviral LTRs (including the modified retroviral LTRs hereinabove described); the β -actin promoter; and human growth hormone promoters. The promoter also may be the native promoter which controls the genes encoding the polypeptides.

The retroviral plasmid vector is employed to transduce packaging cell lines to form producer cell lines. Examples of packaging cells which may be transfected include, but are not limited to, the PE501, PA317, ψ -2, ψ -AM, PA12, T19-14X, VT-19-17-H2, ψ CRE, ψ CRIP, GP+E-86, GP+envAml2, and DAN cell lines as described in Miller, Human Gene Therapy, Vol. 1, pgs. 5-14 (1990), which is incorporated herein by reference.

in its entirety. The vector may transduce the packaging cells through any means known in the art. Such means include, but are not limited to, electroporation, the use of liposomes, and CaPO₄ precipitation. In one alternative, the retroviral plasmid vector may be encapsulated into a liposome, or coupled to a lipid, and then administered to a host. ✓

The producer cell line generates infectious retroviral vector particles which include the nucleic acid sequence(s) encoding the polypeptides. Such retroviral vector particles then may be employed, to transduce eukaryotic cells, either *in vitro* or *in vivo*. The transduced eukaryotic cells will express the nucleic acid sequence(s) encoding the polypeptide. Eukaryotic cells which may be transduced include, but are not limited to, embryonic stem cells, embryonic carcinoma cells, as well as hematopoietic stem cells, hepatocytes, fibroblasts, myoblasts, keratinocytes, endothelial cells, and bronchial epithelial cells. ✓

The present invention also provides a method for determining whether a ligand not known to be capable of binding to a G-protein chemokine receptor can bind to such receptor which comprises contacting a mammalian cell which expresses a G-protein chemokine receptor with the ligand under conditions permitting binding of ligands to the G-protein chemokine receptor, detecting the presence of a ligand which binds to the receptor and thereby determining whether the ligand binds to the G-protein chemokine receptor. The systems hereinabove described for determining agonists and/or antagonists may also be employed for determining ligands which bind to the receptor. ✓

This invention also provides a method of detecting expression of a G-protein chemokine receptor polypeptide of the present invention on the surface of a cell by detecting the presence of mRNA coding for the receptor which comprises obtaining total mRNA from the cell and contacting the mRNA so

obtained with a nucleic acid probe comprising a nucleic acid molecule of at least 10 nucleotides capable of specifically hybridizing with a sequence included within the sequence of a nucleic acid molecule encoding the receptor under hybridizing conditions, detecting the presence of mRNA hybridized to the probe, and thereby detecting the expression of the receptor by the cell. ✓

The present invention also provides a method for identifying receptors related to the receptor polypeptides of the present invention. These related receptors may be identified by homology to a G-protein chemokine receptor polypeptide of the present invention, by low stringency cross hybridization, or by identifying receptors that interact with related natural or synthetic ligands and or elicit similar behaviors after genetic or pharmacological blockade of the chemokine receptor polypeptides of the present invention. ✓

Fragments of the genes may be used as a hybridization probe for a cDNA library to isolate other genes which have a high sequence similarity to the genes of the present invention, or which have similar biological activity. Probes of this type are at least 20 bases, preferably at least 30 bases and most preferably at least 50 bases or more. The probe may also be used to identify a cDNA clone corresponding to a full length transcript and a genomic clone or clones that contain the complete gene of the present invention including regulatory and promoter regions, exons and introns. An example of a screen of this type comprises isolating the coding region of the gene by using the known DNA sequence to synthesize an oligonucleotide probe. Labeled oligonucleotides having a sequence complementary to that of the genes of the present invention are used to screen a library of human cDNA, genomic DNA or mRNA to determine which members of the library the probe hybridizes to. ✓

The present invention also contemplates the use of the genes of the present invention as a diagnostic, for example,

some diseases result from inherited defective genes. These genes can be detected by comparing the sequences of the defective gene with that of a normal one. Subsequently, one can verify that a "mutant" gene is associated with abnormal receptor activity. In addition, one can insert mutant receptor genes into a suitable vector for expression in a functional assay system (e.g., colorimetric assay, expression on MacConkey plates, complementation experiments, in a receptor deficient strain of HBK293 cells) as yet another means to verify or identify mutations. Once "mutant" genes have been identified, one can then screen population for carriers of the "mutant" receptor gene.

Individuals carrying mutations in the gene of the present invention may be detected at the DNA level by a variety of techniques. Nucleic acids used for diagnosis may be obtained from a patient's cells, including but not limited to such as from blood, urine, saliva, tissue biopsy and autopsy material. The genomic DNA may be used directly for detection or may be amplified enzymatically by using PCR (Saiki, et al., Nature, 324:163-166 1986) prior to analysis. RNA or cDNA may also be used for the same purpose. As an example, PCR primers complimentary to the nucleic acid of the instant invention can be used to identify and analyze mutations in the gene of the present invention. For example, deletions and insertions can be detected by a change in size of the amplified product in comparison to the normal genotype. Point mutations can be identified by hybridizing amplified DNA to radio labeled RNA of the invention or alternatively, radio labeled antisense DNA sequences of the invention. Perfectly matched sequences can be distinguished from mismatched duplexes by RNase A digestion or by differences in melting temperatures. Such a diagnostic would be particularly useful for prenatal or even neonatal testing.

Sequence differences between the reference gene and "mutants" may be revealed by the direct DNA sequencing

method. In addition, cloned DNA segments may be used as probes to detect specific DNA segments. The sensitivity of this method is greatly enhanced when combined with PCR. For example, a sequence primer is used with double stranded PCR product or a single stranded template molecule generated by a modified PCR. The sequence determination is performed by conventional procedures with radio labeled nucleotide or by an automatic sequencing procedure with fluorescent-tags. ✓

Genetic testing based on DNA sequence differences may be achieved by detection of alterations in the electrophoretic mobility of DNA fragments in gels with or without denaturing agents. Sequences changes at specific locations may also be revealed by nucleus protection assays, such as RNase and S1 protection or the chemical cleavage method (e.g. Cotton, et al., PNAS, USA, 85:4397-4401 1985). ✓

In addition, some diseases are a result of, or are characterized by changes in gene expression which can be detected by changes in the mRNA. Alternatively, the genes of the present invention can be used as a reference to identify individuals expressing a decrease of functions associated with receptors of this type. ✓

The present invention also relates to a diagnostic assay for detecting altered levels of soluble forms of the G-protein chemokine receptor polypeptides of the present invention in various tissues. Assays used to detect levels of the soluble receptor polypeptides in a sample derived from a host are well known to those of skill in the art and include radioimmunoassays, competitive-binding assays, Western blot analysis and preferably as ELISA assay. ✓

An ELISA assay initially comprises preparing an antibody specific to antigens of the G-protein chemokine receptor polypeptides, preferably a monoclonal antibody. In addition a reporter antibody is prepared against the monoclonal antibody. To the reporter antibody is attached a detectable reagent such as radioactivity, fluorescence or in this

example a horseradish peroxidase enzyme. A sample is now removed from a host and incubated on a solid support, e.g. a polystyrene dish, that binds the proteins in the sample. Any free protein binding sites on the dish are then covered by incubating with a non-specific protein such as bovine serum albumin. Next, the monoclonal antibody is incubated in the dish during which time the monoclonal antibodies attach to any G-protein chemokine receptor proteins attached to the polystyrene dish. All unbound monoclonal antibody is washed out with buffer. The reporter antibody linked to horseradish peroxidase is now placed in the dish resulting in binding of the reporter antibody to any monoclonal antibody bound to G-protein chemokine receptor proteins. Unattached reporter antibody is then washed out. Peroxidase substrates are then added to the dish and the amount of color developed in a given time period is a measurement of the amount of G-protein chemokine receptor proteins present in a given volume of patient sample when compared against a standard curve.

The sequences of the present invention are also valuable for chromosome identification. The sequence is specifically targeted to and can hybridize with a particular location on an individual human chromosome. Moreover, there is a current need for identifying particular sites on the chromosome. Few chromosome marking reagents based on actual sequence data (repeat polymorphisms) are presently available for marking chromosomal location. The mapping of DNAs to chromosomes according to the present invention is an important first step in correlating those sequences with genes associated with disease.

Briefly, sequences can be mapped to chromosomes by preparing PCR primers (preferably 15-25 bp) from the cDNA. Computer analysis of the cDNA is used to rapidly select primers that do not span more than one exon in the genomic DNA, thus complicating the amplification process. These primers are then used for PCR screening of somatic cell

hybrids containing individual human chromosomes. Only those hybrids containing the human gene corresponding to the primer will yield an amplified fragment.

PCR mapping of somatic cell hybrids is a rapid procedure for assigning a particular DNA to a particular chromosome. Using the present invention with the same oligonucleotide primers, sublocalization can be achieved with panels of fragments from specific chromosomes or pools of large genomic clones in an analogous manner. Other mapping strategies that can similarly be used to map to its chromosome include *in situ* hybridization, prescreening with labeled flow-sorted chromosomes and preselection by hybridization to construct chromosome specific-cDNA libraries. ✓

Fluorescence *in situ* hybridization (FISH) of a cDNA clone to a metaphase chromosomal spread can be used to provide a precise chromosomal location in one step. This technique can be used with cDNA as short as 50 or 60 bases. For a review of this technique, see Verma et al., *Human Chromosomes: a Manual of Basic Techniques*, Pergamon Press, New York (1988). ✓

Once a sequence has been mapped to a precise chromosomal location, the physical position of the sequence on the chromosome can be correlated with genetic map data. Such data are found, for example, in V. McKusick, *Mendelian Inheritance in Man* (available on line through Johns Hopkins University Welch Medical Library). The relationship between genes and diseases that have been mapped to the same chromosomal region are then identified through linkage analysis (coinheritance of physically adjacent genes). ✓

Next, it is necessary to determine the differences in the cDNA or genomic sequence between affected and unaffected individuals. If a mutation is observed in some or all of the affected individuals but not in any normal individuals, then the mutation is likely to be the causative agent of the disease. ✓

With current resolution of physical mapping and genetic mapping techniques, a cDNA precisely localized to a chromosomal region associated with the disease could be one of between 50 and 500 potential causative genes. (This assumes 1 megabase mapping resolution and one gene per 20 kb).

The polypeptides, their fragments or other derivatives, or analogs thereof, or cells expressing them can be used as an immunogen to produce antibodies thereto. These antibodies can be, for example, polyclonal or monoclonal antibodies. The present invention also includes chimeric, single chain, and humanized antibodies, as well as Fab fragments, or the product of an Fab expression library. Various procedures known in the art may be used for the production of such antibodies and fragments.

Antibodies generated against the polypeptides corresponding to a sequence of the present invention can be obtained by direct injection of the polypeptides into an animal or by administering the polypeptides to an animal, preferably a nonhuman. The antibody so obtained will then bind the polypeptides itself. In this manner, even a sequence encoding only a fragment of the polypeptides can be used to generate antibodies binding the whole native polypeptides. Such antibodies can then be used to isolate the polypeptide from tissue expressing that polypeptide.

For preparation of monoclonal antibodies, any technique which provides antibodies produced by continuous cell line cultures can be used. Examples include the hybridoma technique (Kohler and Milstein, 1975, Nature, 256:495-497), the trioma technique, the human B-cell hybridoma technique (Kozbor et al., 1983, Immunology Today 4:72), and the EBV-hybridoma technique to produce human monoclonal antibodies (Cole, et al., 1985, in Monoclonal Antibodies and Cancer Therapy, Alan R. Liss, Inc., pp. 77-96).

Techniques described for the production of single chain antibodies (U.S. Patent 4,946,778) can be adapted to produce single chain antibodies to immunogenic polypeptide products of this invention. Also, transgenic mice may be used to express humanized antibodies to immunogenic polypeptide products of this invention. ✓

The present invention will be further described with reference to the following examples; however, it is to be understood that the present invention is not limited to such examples. All parts or amounts, unless otherwise specified, are by weight. ✓

In order to facilitate understanding of the following examples certain frequently occurring methods and/or terms will be described.

"Plasmids" are designated by a lower case p preceded and/or followed by capital letters and/or numbers. The starting plasmids herein are either commercially available, publicly available on an unrestricted basis, or can be constructed from available plasmids in accord with published procedures. In addition, equivalent plasmids to those described are known in the art and will be apparent to the ordinarily skilled artisan. ✓

"Digestion" of DNA refers to catalytic cleavage of the DNA with a restriction enzyme that acts only at certain sequences in the DNA. The various restriction enzymes used herein are commercially available and their reaction conditions, cofactors and other requirements were used as would be known to the ordinarily skilled artisan. For analytical purposes, typically 1 μ g of plasmid or DNA fragment is used with about 2 units of enzyme in about 20 μ l of buffer solution. For the purpose of isolating DNA fragments for plasmid construction, typically 5 to 50 μ g of DNA are digested with 20 to 250 units of enzyme in a larger volume. Appropriate buffers and substrate amounts for particular restriction enzymes are specified by the ✓

manufacturer. Incubation times of about 1 hour at 37°C are ordinarily used, but may vary in accordance with the supplier's instructions. After digestion the reaction is electrophoresed directly on a polyacrylamide gel to isolate the desired fragment.

Size separation of the cleaved fragments is performed using 8 percent polyacrylamide gel described by Goeddel, D. et al., *Nucleic Acids Res.*, 8:4057 (1980).

"Oligonucleotides" refers to either a single stranded polydeoxynucleotide or two complementary polydeoxynucleotide strands which may be chemically synthesized. Such synthetic oligonucleotides have no 5' phosphate and thus will not ligate to another oligonucleotide without adding a phosphate with an ATP in the presence of a kinase. A synthetic oligonucleotide will ligate to a fragment that has not been dephosphorylated.

"Ligation" refers to the process of forming phosphodiester bonds between two double stranded nucleic acid fragments (Maniatis, T., et al., *Id.*, p. 146). Unless otherwise provided, ligation may be accomplished using known buffers and conditions with 10 units to T4 DNA ligase ("ligase") per 0.5 µg of approximately equimolar amounts of the DNA fragments to be ligated.

Unless otherwise stated, transformation was performed as described in the method of Graham, F. and Van der Eb, A., *Virology*, 52:456-457 (1973).

Example 1

Bacterial Expression and Purification of HDGNR10

The DNA sequence encoding for HDGNR10, ATCC # is initially amplified using PCR oligonucleotide primers corresponding to the 5' and sequences of the processed HDGNR10 protein (minus the signal peptide sequence) and the vector sequences 3' to the HDGNR10 gene. Additional nucleotides corresponding to HDGNR10 were added to the 5' and 3' sequences respectively. The 5' oligonucleotide primer has

the sequence 5' CGGAATTCCTCCATGGATTATCAAGTGTCA 3' contains an EcoRI restriction enzyme site followed by 18 nucleotides of HDG NR10 coding sequence starting from the presumed terminal amino acid of the processed protein codon. The 3' sequence 5' CGGAAGCTTCGTCACAAGCCCCACAGATAT 3' contains complementary sequences to a HindIII site and is followed by 18 nucleotides of HDG NR10 coding sequence. The restriction enzyme sites correspond to the restriction enzyme sites on the bacterial expression vector pQE-9 (Qiagen, Inc. 9259 Eton Avenue, Chatsworth, CA, 91311). pQE-9 encodes antibiotic resistance (Amp^r), a bacterial origin of replication (ori), an IPTG-regulatable promoter operator (P/O), a ribosome binding site (RBS), a 6-His tag and restriction enzyme sites. pQE-9 was then digested with EcoRI and HindIII. The amplified sequences were ligated into pQE-9 and were inserted in frame with the sequence encoding for the histidine tag and the RBS. The ligation mixture was then used to transform *E. coli* strain M15/rep 4 (Qiagen, Inc.) by the procedure described in Sambrook, J. et al., Molecular Cloning: A Laboratory Manual, Cold Spring Laboratory Press, (1989). M15/rep4 contains multiple copies of the plasmid pREP4, which expresses the lacI repressor and also confers kanamycin resistance (Kan^r). Transformants are identified by their ability to grow on LB plates and ampicillin/kanamycin resistant colonies were selected. Plasmid DNA was isolated and confirmed by restriction analysis. Clones containing the desired constructs were grown overnight (O/N) in liquid culture in LB media supplemented with both Amp (100 ug/ml) and Kan (25 ug/ml). The O/N culture is used to inoculate a large culture at a ratio of 1:100 to 1:250. The cells were grown to an optical density 600 (O.D.₆₀₀) of between 0.4 and 0.6. IPTG ("Isopropyl-B-D-thiogalacto pyranoside") was then added to a final concentration of 1 mM. IPTG induces by inactivating the lacI repressor, clearing the P/O leading to increased gene expression. Cells were grown an extra 3 to 4 hours.

Cells were then harvested by centrifugation. The cell pellet was solubilized in the chaotropic agent 6 Molar Guanidine HCl. After clarification, solubilized HDGNR10 was purified from this solution by chromatography on a Nickel-Chelate column under conditions that allow for tight binding by proteins containing the 6-His tag. Hochuli, E. et al., J. Chromatography 411:177-184 (1984). HDGNR10 was eluted from the column in 6 molar guanidine HCl pH 5.0 and for the purpose of renaturation adjusted to 3 molar guanidine HCl, 100mM sodium phosphate, 10 mmolar glutathione (reduced) and 2 mmolar glutathione (oxidized). After incubation in this solution for 12 hours the protein was dialyzed to 10 mmolar sodium phosphate. ✓

Example 2

Expression of Recombinant HDGNR10 in COS cells

The expression of plasmid, HDGNR10 HA is derived from a vector pCDNA1/Amp (Invitrogen) containing: 1) SV40 origin of replication, 2) ampicillin resistance gene, 3) E.coli replication origin, 4) CMV promoter followed by a polylinker region, a SV40 intron and polyadenylation site. A DNA fragment encoding the entire HDGNR10 precursor and a HA tag fused in frame to its 3' end was cloned into the polylinker region of the vector, therefore, the recombinant protein expression is directed under the CMV promoter. The HA tag correspond to an epitope derived from the influenza hemagglutinin protein as previously described (I. Wilson, H. Niman, R. Heighten, A. Cherenson, M. Connolly, and R. Lerner, 1984, Cell 37, 767). The infusion of HA tag to the target protein allows easy detection of the recombinant protein with an antibody that recognizes the HA epitope.

The plasmid construction strategy is described as follows: #4

The DNA sequence encoding for HDGNR10, ATCC # was constructed by PCR using two primers: the 5' primer 5' GTCC

AAGCTTGCCACCATGGATTATCAAGTGTC 3' and contains a HindIII site followed by 18 nucleotides of HDG NR10 coding sequence starting from the initiation codon; the 3' sequence 5' CTAGCTCGAGTCAAGCGTAGTCTGGGACGTCGTATGGGTAGCACAAAGCCCCACAGATATTTTC 3' contains complementary sequences to an XhoI site, translation stop codon, HA tag and the last 18 nucleotides of the HDG NR10 coding sequence (not including the stop codon). Therefore, the PCR product contains a HindIII site HDG NR10 coding sequence followed by HA tag fused in frame, a translation termination stop codon next to the HA tag, and an XhoI site. The PCR amplified DNA fragment and the vector, pCDNAI/Amp, were digested with HindIII and XhoI restriction enzyme and ligated. The ligation mixture was transformed into E. coli strain SURE (available from Stratagene Cloning Systems, 11099 North Torrey Pines Road, La Jolla, CA 92037) the transformed culture was plated on ampicillin media plates and resistant colonies were selected. Plasmid DNA was isolated from transformants and examined by restriction analysis for the presence of the correct fragment. For expression of the recombinant HDG NR10, COS cells were transfected with the expression vector by DEAE-DEXTRAN method. (J. Sambrook, E. Fritsch, T. Maniatis, Molecular Cloning: A Laboratory Manual, Cold Spring Laboratory Press, (1989)). The expression of the HDG NR10 HA protein was detected by radiolabelling and immunoprecipitation method. (E. Harlow, D. Lane, Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory Press, (1988)). Cells were labelled for 8 hours with ³⁵S-cysteine two days post transfection. Culture media were then collected and cells were lysed with detergent (RIPA buffer (150 mM NaCl, 1% NP-40, 0.1% SDS, 1% NP-40, 0.5% DOC, 50mM Tris, pH 7.5). (Wilson, I. et al., Id. 37:767 (1984)). Both cell lysate and culture media were precipitated with a HA specific monoclonal antibody. Proteins precipitated were analyzed on 15% SDS-PAGE gels.

Example 3Cloning and expression of HDGNR10 using the baculovirus expression system 5
#

The DNA sequence encoding the full length HDGNR10 protein, ATCC # , was amplified using PCR oligonucleotide primers corresponding to the 5' and 3' sequences of the gene:

The 5' primer has the sequence 5' CGGGATCCCTCCATGGATTAT CAAGTGTCA 3' and contains a BamHI restriction enzyme site followed by 4 nucleotides resembling an efficient signal for the initiation of translation in eukaryotic cells (J. Mol. Biol. 1987, 196, 947-950, Kozak, M.), and just behind the first 18 nucleotides of the HDGNR10 gene (the initiation codon for translation is "ATG").

The 3' primer has the sequence 5' CGGGATCCCGCT CACAAGCCCACAGATAT 3' and contains the cleavage site for the restriction endonuclease BamHI and 18 nucleotides complementary to the 3' non-translated sequence of the HDGNR10 gene. The amplified sequences were isolated from a 1% agarose gel using a commercially available kit ("Geneclean," BIO 101 Inc., La Jolla, Ca.). The fragment was then digested with the endonuclease BamHI and purified as described above. This fragment is designated F2.

The vector pRG1 (modification of pVL941 vector, discussed below) is used for the expression of the HDGNR10 protein using the baculovirus expression system (for review see: Summers, M.D. and Smith, G.E. 1987, A manual of methods for baculovirus vectors and insect cell culture procedures, Texas Agricultural Experimental Station Bulletin No. 1555). This expression vector contains the strong polyhedrin promoter of the Autographa californica nuclear polyhedrosis virus (AcMNPV) followed by the recognition sites for the restriction endonuclease BamHI. The polyadenylation site of the simian virus (SV)40 is used for efficient polyadenylation. For an easy selection of recombinant

viruses the beta-galactosidase gene from *E. coli* is inserted in the same orientation as the polyhedrin promoter followed by the polyadenylation signal of the polyhedrin gene. The polyhedrin sequences are flanked at both sides by viral sequences for the cell-mediated homologous recombination of co-transfected wild-type viral DNA. Many other baculovirus vectors could be used in place of pRG1 such as pAc373, pVL941 and pAcIM1 (Luckow, V.A. and Summers, M.D., *Virology*, 170:31-39).

The plasmid was digested with the restriction enzyme BamHI and then dephosphorylated using calf intestinal phosphatase by procedures known in the art. The DNA was then isolated from a 1% agarose gel as described above. This vector DNA is designated V2.

Fragment F2 and the dephosphorylated plasmid V2 were ligated with T4 DNA ligase. *E. coli* HB101 cells were then transformed and bacteria identified that contained the plasmid (pBachHDGNR10) with the HDGNR10 gene using the enzyme BamHI. The sequence of the cloned fragment was confirmed by DNA sequencing.

5 μ g of the plasmid pBachHDGNR10 were co-transfected with 1.0 μ g of a commercially available linearized baculovirus ("BaculoGold™ baculovirus DNA", Pharmingen, San Diego, CA.) using the lipofection method (Felgner et al. *Proc. Natl. Acad. Sci. USA*, 84:7413-7417 (1987)).

1 μ g of BaculoGold™ virus DNA and 5 μ g of the plasmid pBachHDGNR10 were mixed in a sterile well of a microtiter plate containing 50 μ l of serum free Grace's medium (Life Technologies Inc., Gaithersburg, MD). Afterwards 10 μ l Lipofectin plus 90 μ l Grace's medium were added, mixed and incubated for 15 minutes at room temperature. Then the transfection mixture was added drop wise to the Sf9 insect cells (ATCC CRL 1711) seeded in a 35 mm tissue culture plate with 1 ml Grace' medium without serum. The plate was rocked back and forth to mix the newly added solution. The plate

was then incubated for 5 hours at 27°C. After 5 hours the transfection solution was removed from the plate and 1 ml of Grace's insect medium supplemented with 10% fetal calf serum was added. The plate was put back into an incubator and cultivation continued at 27°C for four days.

After four days the supernatant was collected and a plaque assay performed similar as described by Summers and Smith (supra). As a modification an agarose gel with "Blue Gal" (Life Technologies Inc., Gaithersburg) was used which allows an easy isolation of blue stained plaques. (A detailed description of a "plaque assay" can also be found in the user's guide for insect cell culture and baculovirology distributed by Life Technologies Inc., Gaithersburg, page 9-10).

Four days after the serial dilution, the viruses were added to the cells, blue stained plaques were picked with the tip of an Eppendorf pipette. The agar containing the recombinant viruses was then resuspended in an Eppendorf tube containing 200 µl of Grace's medium. The agar was removed by a brief centrifugation and the supernatant containing the recombinant baculoviruses was used to infect Sf9 cells seeded in 35 mm dishes. Four days later the supernatants of these culture dishes were harvested and then stored at 4°C.

Sf9 cells were grown in Grace's medium supplemented with 10% heat-inactivated FBS. The cells were infected with the recombinant baculovirus V-HDGNR10 at a multiplicity of infection (MOI) of 2. Six hours later the medium was removed and replaced with SF900 II medium minus methionine and cysteine (Life Technologies Inc., Gaithersburg). 42 hours later 5 µCi of ³⁵S-methionine and 5 µCi ³⁵S cysteine (Amersham) were added. The cells were further incubated for 16 hours before they were harvested by centrifugation and the labelled proteins visualized by SDS-PAGE and autoradiography.

Example 4

Expression via Gene Therapy

Fibroblasts are obtained from a subject by skin biopsy. The resulting tissue is placed in tissue-culture medium and separated into small pieces. Small chunks of the tissue are placed on a wet surface of a tissue culture flask, approximately ten pieces are placed in each flask. The flask is turned upside down, closed tight and left at room temperature over night. After 24 hours at room temperature, the flask is inverted and the chunks of tissue remain fixed to the bottom of the flask and fresh media (e.g., Ham's F12 media, with 10% FBS, penicillin and streptomycin, is added. This is then incubated at 37°C for approximately one week. At this time, fresh media is added and subsequently changed every several days. After an additional two weeks in culture, a monolayer of fibroblasts emerge. The monolayer is trypsinized and scaled into larger flasks.

pMV-7 (Kirschmeier, P.T. et al, DNA, 7:219-25 (1988)) flanked by the long terminal repeats of the Moloney murine sarcoma virus, is digested with EcoRI and HindIII and subsequently treated with calf intestinal phosphatase. The linear vector is fractionated on agarose gel and purified, using glass beads.

The cDNA encoding a polypeptide of the present invention is amplified using PCR primers which correspond to the 5' and 3' end sequences respectively. The 5' primer contains an EcoRI site and the 3' primer contains a HindIII site. Equal quantities of the Moloney murine sarcoma virus linear backbone and the EcoRI and HindIII fragment are added together, in the presence of T4 DNA ligase. The resulting mixture is maintained under conditions appropriate for ligation of the two fragments. The ligation mixture is used to transform bacteria HB101, which are then plated onto agar-containing kanamycin for the purpose of confirming that the vector had the gene of interest properly inserted.

The amphotropic pA317 or GP+aml2 packaging cells are grown in tissue culture to confluent density in Dulbecco's Modified Eagles Medium (DMEM) with 10% calf serum (CS), penicillin and streptomycin. The MSV vector containing the gene is then added to the media and the packaging cells are transduced with the vector. The packaging cells now produce infectious viral particles containing the gene (the packaging cells are now referred to as producer cells).

Fresh media is added to the transduced producer cells, and subsequently, the media is harvested from a 10 cm plate of confluent producer cells. The spent media, containing the infectious viral particles, is filtered through a millipore filter to remove detached producer cells and this media is then used to infect fibroblast cells. Media is removed from a sub-confluent plate of fibroblasts and quickly replaced with the media from the producer cells. This media is removed and replaced with fresh media. If the titer of virus is high, then virtually all fibroblasts will be infected and no selection is required. If the titer is very low, then it is necessary to use a retroviral vector that has a selectable marker, such as neo or his.

The engineered fibroblasts are then injected into the host, either alone or after having been grown to confluence on cytodex 3 microcarrier beads. The fibroblasts now produce the protein product.

Numerous modifications and variations of the present invention are possible in light of the above teachings and, therefore, within the scope of the appended claims, the invention may be practiced otherwise than as particularly described.

SEQUENCE LISTING

- (1) GENERAL INFORMATION:
- (i) APPLICANT: Li, ET AL.
- (ii) TITLE OF INVENTION: Human G-Protein Chemokine Receptor
- (iii) NUMBER OF SEQUENCES:
- (iv) CORRESPONDENCE ADDRESS:
- (A) ADDRESSEE: CARELLA, BYRNE, BAIN, GILFILLAN, CECCHI, STEWART & OLSTEIN
- (B) STREET: 6 BECKER FARM ROAD
- (C) CITY: ROSELAND
- (D) STATE: NEW JERSEY
- (E) COUNTRY: USA
- (F) ZIP: 07068
- (v) COMPUTER READABLE FORM:
- (A) MEDIUM TYPE: 3.5 INCH DISKETTE
- (B) COMPUTER: IBM PS/2
- (C) OPERATING SYSTEM: MS-DOS
- (D) SOFTWARE: WORD PERFECT 5.1
- (vi) CURRENT APPLICATION DATA:
- (A) APPLICATION NUMBER:
- (B) FILING DATE: concurrently
- (C) CLASSIFICATION:
- (vii) ATTORNEY/AGENT INFORMATION:
- (A) NAME: FERRARO, GREGORY D.
- (B) REGISTRATION NUMBER: 36,134
- (C) REFERENCE/DOCKET NUMBER: 325800-
- (viii) TELECOMMUNICATION INFORMATION:
- (A) TELEPHONE: 201-994-1700
- (B) TELEFAX: 201-994-1744
- (2) INFORMATION FOR SEQ ID NO:1:
- (i) SEQUENCE CHARACTERISTICS
- (A) LENGTH: 1414 BASE PAIRS
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: SINGLE
- (D) TOPOLOGY: LINEAR
- (ii) MOLECULE TYPE: cDNA
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

GTGAGATGGT GCTTTCATGA ATTCCCTCAA CAAGAGCCAA GCTCTCCATC TAGTGGACAG 60

GGAAGCTAGC AGCAAACCTT CCCTTCACTA CGAAACTTCA TTGCTTGGCC CAAAAGAGAG 120
 TTAATTCAAT GTAGACATCT ATGTAGGCAA TTA AAAACCT ATTGATGTAT AAAACAGTTT 180
 GCATTTCATGG AGGGCACTA AATACATTCT AGGACTTTAT AAAAGATCAC TTTTATTTA 240
 TGCACAGGGT GGAACAAG ATG GAT TAT CAA GTG TCA AGT CCA ATC TAT GAC 291
 Met Asp Tyr Gln Val Ser Ser Pro Ile Tyr Asp
 ATC AAT TAT TAT ACA TCG GAG CCC TGC CCA AAA ATC AAT GTG AAG CAA 339
 Ile Asn Tyr Tyr Thr Ser Glu Pro Cys Pro Lys Ile Asn Val Lys Gln
 ATC GCA GCC CGC CTC CTG CCT CCG CTC TAC TCA CTG GTG TTC ATC TTT 387
 Ile Ala Ala Arg Leu Leu Pro Pro Leu Tyr Ser Leu Val Phe Ile Phe
 GGT TTT GTG GGC AAC ATG CTG GTC ATC CTC ATC CTG ATA AAC TGC CAA 435
 Gly Phe Val Gly Asn Met Leu Val Ile Leu Ile Leu Ile Asn Cys Gln
 AGG CTG GAG AGC ATG ACT GAC ATC TAC CTG CTC AAC CTG GCC ATC TCT 483
 Arg Leu Glu Ser Met Thr Asp Ile Tyr Leu Leu Asn Leu Ala Ile Ser
 GAC CTG TTT TTC CTT CTT ACT GTC CCC TTC TGG GCT CAC TAT GCT GCC 531
 Asp Leu Phe Phe Leu Leu Thr Val Pro Phe Trp Ala His Tyr Ala Ala
 GCC CAG TGG GAC TTT GGA AAT ACA ATG TGT CAA CTC TTG ACA GGG CTC 579
 Ala Gln Trp Asp Phe Gly Asn Thr Met Cys Leu Leu Thr Gly Leu Tyr
 TAT TTT ATA GGC TTC TTC TCT GGA ATC TTC TTC ATC ATC CTC CTG ACA 627
 Phe Ile Gly Phe Phe Ser Gly Ile Phe Phe Ile Ile Gln Leu Leu Thr
 ATC GAT AGG TAC CTG GCT ATC GTC CAT GCT GTG TTT GCT TTA AAA GCC 675
 Ile Asp Arg Tyr Leu Ala Ile Val His Ala Val Phe Ala Leu Lys Ala
 AGG ACG GTC ACC TTT GGG GTG GTG ACA AGT GTG ATC ACT TGG GTG GTG 723
 Arg Thr Val Thr Phe Gly Val Val Thr Ser Val Ile Thr Trp Val Val
 GCT GTG TTT GCG TCT CTC CCA GGA ATC ATC TTT ACC AGA TCT CAA AAA 771
 Ala Val Phe Ala Ser Leu Pro Gly Ile Ile Phe Thr Arg Ser Gln Lys
 GAA GGT CTT CAT TAC ACC TGC AGC TCT CAT TTT CCA TAC AGT CAG TAT 819
 Glu Gly Leu His Tyr Thr Cys Ser Ser His Phe Pro Tyr Ser Gln Tyr
 CAA TTC TGG AAG AAT TTC CAG ACA TTA AAG ATA GTC ATC TTG GGG CTG 867
 Gln Phe Trp Lys Asn Phe Gln-Thr Leu Lys Ile Val Ile Leu Gly Leu
 GTC CTG CCG CTG CTT GTC ATG GTC ATC TGC TAC TCG GGA ATC CTA AAA 915
 Val Leu Pro Leu Leu Val Met Val Ile Cys Tyr Ser Gly Ile Leu Lys
 ACT CTG CTT CGG TGT CGA AAT GAG AAG AAG AGG CAC AGG GCT GTG AGG 963
 Thr Leu Leu Arg Cys Arg Asn Glu Lys Lys Arg His Arg Ala Val Arg
 CTT ATC TTC ACC ATC ATG ATT GTT TAT TTT CTC TTC TGG GCT CCC TAC 1011
 Leu Ile Phe Thr Ile Met Ile Val Tyr Phe Leu Phe Trp Ala Pro Tyr
 AAC ATT GTC CTT CTC CTG AAC ACC TTC CAG GAA TTC TCT GGC CTG AAT 1059
 Asn Ile Val Leu Leu Leu Asn Thr Phe Gln Glu Phe Phe Gly Leu Asn
 AAT TGC AGT AGC TCT AAC AGG TTG GAC CAA GCT ATG CAG GTG ACA GAG 1107
 Asn Cys Ser Ser Ser Asn Arg Leu Asp Gln Ala Met Gln Val Thr Glu
 ACT CTT GGG ATG ACG CAC TGC TGC ATC AAC CCC ATC ATC TAT GCC TTT 1155
 Thr Leu Gly Met Thr His Cys Cys Ile Asn Pro Ile Ile Tyr Ala Phe

GTC GGG GAG AAG TTC AGA AAC TAC CTC TTA GTC TTC TTC CAA AAG CAC 1203
 Val Gly Glu Lys Phe Arg Asn Tyr Leu Leu Val Phe Phe Gln Lys His
 ATT GCC AAA CGC TTC TGC AAA TGC TGT TCT ATT TTC CAG CAA GAG GCT 1251
 Ile Ala Lys Arg Phe Cys Lys Cys Cys Ser Ile Phe Gln Gln Glu Ala
 CCC GAG CGA GCA AGC TCA GTT TAC ACC CGA TCC ACT GGG GAG CAG GAA 1299
 Pro Glu Arg Ala Ser Ser Val Tyr Thr Arg Ser Thr Gly Glu Gln Glu
 ATA TCT GTG GGC TTG TGACACGGAC TCAAGTGGGC TGGTGACCCA GTCAGAGTTG 1354
 Ile Ser Val Gly Leu
 TGCACATGGC TTAGTTTTCA TACACAGCCT GGGCTGGGGG TGGGGTGGAA GAGGTCTTTT 1414

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS

- (A) LENGTH: AMINO ACIDS
 (B) TYPE: AMINO ACID
 (C) STRANDEDNESS:
 (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met	Asp	Tyr	Gln	Val	Ser	Ser	Pro	Ile	Tyr	Asp	Ile	Asn	Tyr	Tyr	5	10	15
Thr	Ser	Glu	Pro	Cys	Pro	Lys	Ile	Asn	Val	Lys	Gln	Ile	Ala	Ala	20	25	30
Arg	Leu	Leu	Pro	Pro	Leu	Tyr	Ser	Leu	Val	Phe	Ile	Phe	Gly	Phe	35	40	45
Val	Gly	Asn	Met	Leu	Val	Ile	Leu	Ile	Leu	Ile	Asn	Cys	Gln	Arg	50	55	60
Leu	Glu	Ser	Met	Thr	Asp	Ile	Tyr	Leu	Leu	Asn	Leu	Ala	Ile	Ser	65	70	75
Asp	Leu	Phe	Phe	Leu	Leu	Thr	Val	Pro	Phe	Trp	Ala	His	Tyr	Ala	80	85	90
Ala	Ala	Gln	Trp	Asp	Phe	Gly	Asn	Thr	Met	Cys	Leu	Leu	Thr	Gly	95	100	105
Leu	Tyr	Phe	Ile	Gly	Phe	Phe	Ser	Gly	Ile	Phe	Phe	Ile	Ile	Gln	110	115	120
Leu	Leu	Thr	Ile	Asp	Arg	Tyr	Leu	Ala	Ile	Val	His	Ala	Val	Phe	125	130	135
Ala	Leu	Lys	Ala	Arg	Thr	Val	Thr	Phe	Gly	Val	Val	Thr	Ser	Val	140	145	150
Ile	Thr	Trp	Val	Val	Ala	Val	Phe	Ala	Ser	Leu	Pro	Gly	Ile	Ile	155	160	165
Phe	Thr	Arg	Ser	Gln	Lys	Glu	Gly	Leu	His	Tyr	Thr	cys	Ser	Ser	170	175	180
His	Phe	Pro	Tyr	Ser	Gln	Tyr	Gln	Phe	Trp	Lys	Asn	Phe	Gln	Thr	185	190	195
Leu	Lys	Ile	Val	Ile	Leu	Gly	Leu	Val	Leu	Pro	Leu	Leu	Val	Met	200	205	210

Val	Ile	Cys	Tyr	Ser	Gly	Ile	Leu	Lys	Thr	Leu	Leu	Arg	Cys	Arg	
				215					220					225	
Asn	Glu	Lys	Lys	Arg	His	Arg	Ala	Val	Arg	Leu	Ile	Phe	Thr	Ile	
				230					235					240	
Met	Ile	Val	Tyr	Phe	Leu	Phe	Trp	Ala	Pro	Tyr	Asn	Ile	Val	Leu	
				245					250					255	
Leu	Leu	Asn	Thr	Phe	Gln	Glu	Phe	Phe	Gly	Leu	Asn	Asn	Cys	Ser	
				260					265					270	
Ser	Ser	Asn	Arg	Leu	Asp	Gln	Ala	Met	Gln	Val	Thr	Glu	Thr	Leu	
				275					280					285	
Gly	Met	Thr	His	Cys	Cys	Ile	Asn	Pro	Ile	Ile	Tyr	Ala	Phe	Val	
				290					295					300	
Gly	Glu	Lys	Phe	Arg	Asn	Tyr	Leu	Leu	Val	Phe	Phe	Gln	Lys	His	
				305					310					315	
Ile	Ala	Lys	Arg	Phe	Cys	Lys	Cys	Cys	Ser	Ile	Phe	Gln	Gln	Glu	
				320					325					330	
Ala	Pro	Glu	Arg	Ala	Ser	Ser	Val	Tyr	Thr	Arg	Ser	Thr	Gly	Glu	
				335					340					345	
Gln	Glu	Ile	Ser	Val	Gly	Leu									
				350											

WHAT IS CLAIMED IS:

1. An isolated polynucleotide comprising a member selected from the group consisting of:
 - (a) a polynucleotide encoding the polypeptide as set forth in SEQ ID NO:2;
 - (b) a polynucleotide encoding a mature polypeptide encoded by the DNA contained in ATCC Deposit No. _____;
 - (c) a polynucleotide capable of hybridizing to and which is at least 70% identical to the polynucleotide of (a) or (b); and
 - (d) a polynucleotide fragment of the polynucleotide of (a), (b) or (c).
2. The polynucleotide of claim 1 wherein the polynucleotide is DNA.
3. A vector containing the DNA of Claim 2.
4. A host cell transformed or transfected with the vector of Claim 3.
5. A process for producing a polypeptide comprising: expressing from the host cell of Claim 4 the polypeptide encoded by said DNA.
6. A process for producing cells capable of expressing a polypeptide comprising transforming or transfecting the cells with the vector of Claim 3.
7. A receptor polypeptide comprising a member selected from the group consisting of:
 - (i) a polypeptide having the deduced amino acid sequence of SEQ ID NO:2 and fragments, analogs and derivatives thereof; and

(ii) a polypeptide encoded by the cDNA of ATCC Deposit No. _____ and fragments, analogs and derivatives of said polypeptide.

8. The polypeptide of Claim 7 wherein the polypeptide has the deduced amino acid sequence of SEQ ID NO:2.
9. An antibody against the polypeptide of claim 7 selected from the group consisting of an antibody which agonizes the activity of the polypeptide and an antibody which antagonizes the activity of the polypeptide.
10. A compound which activates the polypeptide of claim 7.
11. A compound which inhibits activation the polypeptide of claim 7.
12. A method for the treatment of a patient having need to activate a G-protein chemokine receptor comprising: administering to the patient a therapeutically effective amount of the compound of claim 10.
13. A method for the treatment of a patient having need to inhibit a G-protein chemokine receptor comprising: administering to the patient a therapeutically effective amount of the compound of claim 11.
14. The method of claim 12 wherein said compound is a polypeptide and a therapeutically effective amount of the compound is administered by providing to the patient DNA encoding said agonist and expressing said agonist *in vivo*.

15. The method of claim 13 wherein said compound is a polypeptide and a therapeutically effective amount of the compound is administered by providing to the patient DNA encoding said antagonist and expressing said antagonist *in vivo*.
16. A method for identifying compounds which bind to and activate the receptor polypeptide of claim 7 comprising:
contacting a cell expressing on the surface thereof the receptor polypeptide, said receptor being associated with a second component capable of providing a detectable signal in response to the binding of a compound to said receptor polypeptide, with a compound under conditions sufficient to permit binding of the compound to the receptor polypeptide; and
identifying if the compound is an effective agonist by detecting the signal produced by said second component.
17. A method for identifying compounds which bind to and inhibit activation the polypeptide of claim 7 comprising:
contacting a cell expressing on the surface thereof the receptor polypeptide, said receptor being associated with a second component capable of providing a detectable signal in response to the binding of a compound to said receptor polypeptide, with a compound to be screened under conditions to permit binding to the receptor polypeptide; and
determining whether the compound inhibits activation of the polypeptide by detecting the absence of a signal generated from the interaction of the ligand with the polypeptide.

18. A process for diagnosing a disease or a susceptibility to a disease related to an under-expression of the polypeptide of claim 7 comprising:
determining a mutation in the nucleic acid sequence encoding said polypeptide.
19. The polypeptide of Claim 7 wherein the polypeptide is a soluble fragment of the polypeptide and is capable of binding a ligand for the receptor.
20. A diagnostic process comprising:
analyzing for the presence of the polypeptide of claim 19 in a sample derived from a host.

FIG. 1A

10 GTGAGATGGTGTCTTTCATGAATTCCCCCAACAAGAGCCCAAGCTCTCCATCTAGTGGACAG 50
 70 GGAAGCTAGCAGCAACCTTCCCTTCACTACGAACTTCATTGCTTGGCCCAAAAGAGAG 110
 130 TTAATTCAATGTAGACATCTATGTAGGCAATTAAAAACCTATTGATGTATAAAACAGTTT 170
 190 GCATTATGGAGGGCAACTAAATACATTCTAGGACTTTTATAAAAGATCACTTTTATTATA 230
 250 TGCACAGGTGGAACAAGATGGATTATCAAGTGTCAAGTCCAATCTATGACATCAATTAT 290
 310 M D Y Q V S S P I Y D I N Y
 TATACATCGGAGCCCTGCCCAAAATCAATGTGAAGCAAAATCGCAGCCCGCCTCCTGCCT 350
 370 Y T S E P C P K I N V K Q I A A R L L P
 CCGCTCTACTCACTGGTGTTCATCTTTGGTTTGTGGGCAACATGCTGGTCATCCCTCATC 410
 430 P L Y S L V F I F G F V G N M L V I L I
 CTGATAAACTGCCAAAGGCTGGAGAGCATGACTGACATCTACCTGCTCAACCTGGCCATC 470
 490 L I N C Q R L E S M T D I Y L L N L A I
 TCTGACCTGTTTTCCTTCTTACTGTCCCTTCTGGGCTCACTATGCTGCCGCCAGTGG 530
 550 S D L F F L L T V P F W A H Y A A Q W
 GACTTTGGAATACAAATGTGTCAACTCTTGACAGGGCTCTATTTTATAGGCTTCTCTCT 590
 D F G N T M C Q L L T G L Y F I G F F S
 MATCH WITH FIG. 1B

SUBSTITUTE SHEET (RULE 26)

FIG. 1B

MATCH WITH FIG. 1A: 610 630
 GGAATCTTCTTCATCATCCTCCTGACAATCGATAGGTACCTGGCTATCGTCCATGCTGTG 650
 G I F I I L L T I D R Y L A I V H A V
 670 690 710
 TTTCGCTTTAAAGCCAGGACGGTCACCTTTGGGTGGTGACAAGTGTGATCACTTGGGTG
 F A L K A R T V T F G V V T S V I T W V
 730 750 770
 GTGGCTGTGTTGCGTCTCTCCAGGAATCATCTTTACCAGATCTCAAAAAGAGGTCTT
 V A V F A S L P G I I F T R S Q K E G L
 790 810 830
 CATTACACCTGCAGCTCTCATTTTCCATACAGTCAGTATCAATTCTGGAAGAATTTCAG
 H Y T C S S H F P Y S Q Y Q F W K N F Q
 850 870 890
 ACATTAAAGATAGTCATCTTGGGCTGGTCCCTGCCGCTGTTGTTCATGGTCATCTGCTAC
 T L K I V I L G L V L P L L V M V I C Y
 910 930 950
 TCGGGAATCCTAAAACTCTGCTTCGGTTCGAAATGAGAAGAGGCACAGGGCTGTG
 S G I L K T L L R C R N E K K R H R A V
 970 990 1010
 AGGCTTATCTTCACCATCATGATTGTTTATTTTCTCTCTGCGCTCCCTACAACATTGTC
 R L I F T I M I V Y F L F W A P Y N I V
 1030 1050 1070
 MATCH WITH FIG. 1C

SUBSTITUTE SHEET (RULE 26)

MATCH WITH FIG. 1B

FIG. 1C

CTTCTCCTGAACACCTTCCAGGAATCTTTGGCCTGAATAATGCGTAGCTCTAACAGG
L L L N T F Q E F F G L N N C S S S N R
1090
TTGGACCAAGCTATGCAGGTGACAGAGACTCTTGGGATGACGCACCTGCTGCATCAACCCC
L D Q A M Q V T E T L G M T H C C I N P
1110
ATCATCTATGCCCTTTGTCTGGGAGAGTTCAGAACTACCTCTTAGTCTTCTCCAAAAG
I I Y A F V G E K F R N Y L L V F F Q K
1130
CACATTGCCAAACGCTTCTGCAAAATGCTGTCTATTTCCAGCAAGAGGCTCCCGAGCGA
H I A K R F C K C C S I F Q Q E A P E R
1210
GCAAGCTCAGTTTACACCCGATCCACTGGGGAGCAGGAAATATCTGTGGCTGTGTGACAC
A S S V Y T R S T G E Q E I S V G L *
1230
GGACTCAAGTGGGCTGGTGACCCAGTCAGAGTTGTGCACATGGCTTAGTTTCATACACA
1250
GCCTGGGCTGGGGTGGGTGGAAGAGGCTTTT
1270
1290
1310
1330
1350
1370
1390
1410

SUBSTITUTE SHEET (RULE 26)

4 / 4

4 QVSSPIYDINYYTSEPCPKINVKQIAARLLPPLYSLVFI FGVGNMLVIL 53
 18 EEVTTFFDYDY..GAPCHKFDVKQIGAQLLPPLYSLVFI FGVGNMLVVL 65
 54 ILINCQRLSEMTDIYLLNLAISDLFFELLTPFWAHYAAQWDFGNTMCQL 103
 66 ILINCKKLKCLTDIYLLNLAISDLLFLITPLWAHSAANEWVFGNAMCKL 115
 104 LTGLYFIGFFSGIFFIILLTIDRYLAIVHAVFALKARTVTFGVVTSVITW 153
 116 FTGLYHIGYFGGIFFIILLTIDRYLAIVHAVFALKARTVTFGVVTSVITW 153
 154 VVAVFASLPGIIFTRSQEGLHYTCSSHFPYSQYQFWKNFQTLKIVILGL 203
 166 LVAVFASVPGIIFTKCQKEDSVVCGPYFPRG...WNNFHTIMRNILGL 211
 204 VLPLLVMVICYSGILKTLRLCRNEKKRHRVRLIFTIMIVYFLFWAPYNI 253
 212 VLPLLIMVICYSGILKTLRLCRNEKKRHRVRLIFTIMIVYFLFWAPYNI 253
 254 VLLNTFQEFFGLNCCSSNRLDQAMQVTTETLGMTHCCINPIIYAFVGEK 261
 262 VILLNTFQEFFGLSNCESTSQLDQATQVTTETLGMTHCCINPIIYAFVGEK 303
 304 FRNYLLVFFQKHIKRFCCKCSIFQQEAPERASSVYTRS...TGEQEISV 350
 312 FRSLFHIALGCRIA.PLQKPVCGGPGVRPGKNVKTQTQGLLDGRGKGKSI 360
 351 G 351
 361 G 361

FIG. 2

SUBSTITUTE SHEET (RULE 26)

Applicant's or agent's file reference number: 325800-369	International application No.: PCT/US95/07173
---	---

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

<p>A. The indications made below relate to the microorganism referred to in the description on pages 6, 34, 36, 37, and (48) lines 43, 11, 11, 19, and (10, 53), respectively.</p>	
<p>B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/></p>	
<p>Name of depository institution</p> <p>American Type Culture Collection</p>	
<p>Address of depository institution (including postal code and country)</p> <p>12301 Parklawn Drive Rockville, Maryland 20852 U.S.A.</p>	
<p>Date of deposit: June 1, 1995</p>	<p>Accession Number: 97183</p>
<p>C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is contained on an additional sheet <input type="checkbox"/></p>	
<p>D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)</p>	
<p>E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)</p> <p>The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")</p>	

For receiving Office use only		For International Bureau use only	
<input type="checkbox"/> This sheet was received with the international application		<input type="checkbox"/> This sheet was received by the International Bureau on:	
Authorized officer		01 NOVEMBER 1996 61.11.96 Authorized officer	

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US95/07173

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : C07K 14/705; C12N 15/12

US CL : 435/6, 69.1, 252.3, 320.1; 530/350; 536/23.5

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/6, 69.1, 252.3, 320.1; 530/350; 536/23.5

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
NONE

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
APS, STN, MEDLINE
search terms: G protein, receptor#, orphan#

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	Proceedings of the National Academy of Science, Vol. 87, issued April 1990, P. C. Ross et al., "RTA, a candidate G protein-coupled receptor: Cloning, sequencing, and tissue distribution", pages 3052-3056, see entire document.	1-8, 16-20
Y	SCIENCE, Vol. 244, issued 05 May 1989, F. Libert et al., "Selective Amplification and Cloning of Four New Members of the G Protein-Coupled Receptor Family", pages 569 to 572, see entire document.	1-8, 16-20
Y	FEBS LETTERS, Vol. 271, issued October 1990, C. Eva et al., "Molecular cloning of a novel G protein-coupled receptor that may belong to the neuropeptide receptor family", pages 81 to 84, see entire document.	1-8, 16-20

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* "A" document defining the general state of the art which is not considered to be of particular relevance	* "T" later documents published after the international filing date or priority date and not in conflict with the application but cited to understand the principles or theory underlying the invention
* "E" earlier document published on or after the international filing date	* "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
* "L" documents which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reasons (as specified)	* "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
* "O" document referring to an oral disclosure, use, exhibition or other means	* "A" document member of the same patent family
* "P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

01 SEPTEMBER 1995

Date of mailing of the international search report

15 SEP 1995

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

JOHN P. ULM

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US95/07173

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>The Journal of Biological Chemistry, Vol. 265, No. 16, issued 05 June 1990, T. Hla et al., "An Abundant Transcript Induced in Differentiating Human Endothelial Cells Encodes a Polypeptide with Structural Similarities to G-protein-coupled Receptors", pages 9306 to 9313, see entire document.</p>	1-8, 16-20

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US95/07173

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:
Please See Extra Sheet.

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.: 1 to 8 and 16 to 20

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US95/07173

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

I. This International Search Authority has found 6 inventions claimed in the International Application covered by the claims indicated below:
Group I, claims 1 to 8 and 16 to 20, drawn to a nucleic acid encoding a receptor polypeptide, the receptor polypeptide encoded thereby and a method of use.

Group II, claim 9, drawn to an antibody which binds to a receptor polypeptide.

Group III, claim 10 drawn to an agonist for a receptor polypeptide.

Group IV, claims 11, drawn to a receptor polypeptide antagonist.

Group V, claims 12 and 14, drawn to a method of use of the product of Group III.

Group VI, claims 13 and 15, drawn to a method of use of the product of Group IV.

and it considers that the International Application does not comply with the requirements of unity of invention (Rules 13.1, 13.2 and 13.3) for the reasons indicated below:

This application contains the following inventions or groups of inventions which are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for all inventions to be examined, the appropriate additional examination fees must be paid.

The inventions listed as Groups I-IV do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons:

Inventions I, II, III and IV are four chemically and structurally different products and as such do not share a common special technical feature. Lack of unity is further shown because each of these inventions can be made and used without any one or more of the other inventions. The inventions of Group V and VI are properly grouped, pursuant to 37 CFR 1.475(d), separately since they do not constitute inventions grouped with the main invention.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☐ BLACK BORDERS

☒ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES

☒ FADED TEXT OR DRAWING

☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING

☒ SKEWED/SLANTED IMAGES

☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS

☐ GRAY SCALE DOCUMENTS

☒ LINES OR MARKS ON ORIGINAL DOCUMENT

☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY

☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.